

Landau-Zener Problem for Trilinear Hamiltonians

Artur Ishkhanyan

*Engineering Center of Armenian National Academy of Sciences,
Ashtarak-2, 378410 Armenia*

Matt Mackie

*Helsinki Institute of Physics, University of Helsinki, PL 64,
FIN-00014 Helsingin yliopisto, Finland*

Andrew Carmichael, Phillip L. Gould, and Juha Javanainen

*Department of Physics, University of Connecticut, Storrs,
Connecticut 06269-3046*

(February 2, 2008)

We consider a nonlinear version of the Landau-Zener problem, focusing on photoassociation of a Bose-Einstein condensate as a specific example. Contrary to the exponential rate dependence obtained for the linear problem, a series expansion technique indicates that, when the resonance is crossed slowly, the probability for failure of adiabaticity is directly proportional to the rate at which the resonance is crossed.

When external conditions force the diabatic energies of two interacting quantum states cross, under conditions of complete adiabaticity the system makes a transition from one bare energy eigenstate to the other. However, “complete adiabaticity” applies when the levels cross “infinitely slowly”. The realistic case when only a finite time is spent near resonance makes the Landau-Zener (LZ) problem [1,2], one of the handful of key paradigms in quantum mechanics. On the other hand, the Gross-Pitaevskii equation and its variants, nonlinear versions of standard quantum mechanics, have proven highly successful in the theoretical description of alkali-vapor Bose-Einstein condensate [3]. Nonlinear quantum mechanics gives up the superposition principle and hence the linear vector space structure. How much of our intuitive understanding of quantum mechanics survives becomes an issue.

The nonlinear version of the LZ problem that is the focus of this Letter is something we have come across in our studies of photoassociation [4,5] of a Bose-Einstein condensate (BEC). However, the same question will come up in an attempt to control the scattering length [6] of an atomic BEC by means of a Feshbach resonance [7], in second-harmonic generation in nonlinear optics [8], and generally in field theories where the nonlinearity is of the form $\psi^\dagger\phi\phi$. For concreteness we couch our discussion in terms of photoassociation, and thus consider processes in which two atoms and a laser photon join to make a molecule. It is known from a second-quantized numerical solution that by sweeping the frequency of the photoassociating laser, in the adiabatic limit it is possible to convert an atomic BEC entirely into a molecular BEC [4]. The question is, what are the nonlinear consequences if the frequency is not swept “infinitely slowly”? While we actually know of no publication addressing this ques-

tion [9,10], we have found that the same applies also to the semiclassical version of the problem, when the atomic and molecular condensates are described by a nonlinear two-state system.

The answer to the above question is outlined as follows. First the essentials of the linear LZ problem are reviewed, which serves both as an introduction to the formalism and a foundation for our series expansion. Turning to the nonlinear LZ model, we use photoassociation as an example to review the dressed state picture and illustrate the physics of adiabatic following. In particular, a picture in terms of crossing of stationary states still applies and predicts near-adiabaticity, but the details are qualitatively different from the case of linear quantum mechanics. The first-order Heisenberg equations of motion for the semi-classical atom-molecule amplitudes are then converted into a third-order differential equation for the molecular probability, whereby a perturbative expansion allows for what amounts to a Green function solution. All told, we find that, while in the linear case the probability for no transition is an exponential function of the speed at which the resonance is crossed, at slow sweep rates of the laser frequency the probability that an atom does not make a transition to the molecular condensate is directly proportional to the rate. Before closing, we estimate explicit numbers for two systems, ^{23}Na and ^{87}Rb , that could serve to test our prediction.

The linear LZ problem is specified by the equations

$$i\dot{\alpha} = \Omega\beta, \quad i\dot{\beta} = \Omega\alpha + \delta\beta. \quad (1)$$

Here α and β are the probability amplitudes for the two bare or diabatic states, Ω is the (real) coupling between them, and the detuning $\delta = \xi t$ stands for the time dependent frequency difference between the two states. For any fixed δ the system (1) has two nonde-

generate dressed [11] or adiabatic states $[\bar{\alpha}, \bar{\beta}]$ with the property that the time dependent solution is of the form $[\alpha(t), \beta(t)] = e^{-i\epsilon t}[\bar{\alpha}, \bar{\beta}_0]$ for some quasienergy $\hbar\epsilon$. In particular, one of these states is such that $[\bar{\alpha}, \bar{\beta}] \rightarrow [1, 0]$ when $\delta \rightarrow -\infty$, and evolves continuously to $[0, 1]$ as $\delta \rightarrow \infty$.

Suppose that the system with the time dependent detuning $\delta = \xi t$ starts out in the state with $\alpha = 1$ when $t \rightarrow -\infty$, then it also start out in this dressed state. As the detuning is swept “infinitely slowly” to $+\infty$, by the adiabatic theorem the system emerges at time $t = \infty$ in the same dressed state, or with the probability amplitudes that satisfy $|\alpha| = 0, |\beta| = 1$. More accurately, when the detuning is not swept infinitely slowly, $\xi > 0$, adiabaticity tends to break down especially in the neighborhood of $\delta = 0$ where the two quasienergies have their closest approach. The time evolution of the probability for state β , $P(t) \equiv |\beta(t)|^2$, often written [2] using parabolic cylinder functions [12], can more conveniently be expressed in terms of the Kummer and Tricomi confluent hypergeometric functions ${}_1F_1$ and U [12]:

$$P_{LZ}(\lambda, t) = 1 - e^{-\pi\lambda} |U(-i\lambda, \frac{1}{2}, -\frac{1}{2}i\xi t^2)|^2, \quad t \leq 0, \quad (2a)$$

$$P_{LZ}(\lambda, t) = 1 - e^{-\pi\lambda} \left| \frac{2\sqrt{\pi}}{\Gamma(\frac{1}{2} - i\lambda)} {}_1F_1(-i\lambda, \frac{1}{2}, -\frac{1}{2}i\xi t^2) + U(-i\lambda, \frac{1}{2}, -\frac{1}{2}i\xi t^2) \right|^2, \quad t \geq 0, \quad (2b)$$

where Γ is the gamma function. At $t = +\infty$ we have the Landau-Zener result

$$P_{LZ}^\infty(\lambda) \equiv P_{LZ}(\lambda, \infty) = 1 - e^{-4\pi\lambda}, \quad (3)$$

where

$$\lambda = \frac{\Omega^2}{2\xi} \quad (4)$$

is the conventional LZ parameter.

Let us next turn to the nonlinear LZ problem, for which we develop a series expansion solution. Modern mathematical physics is of course replete with series expansion techniques. For example, series expansions have been used to solve an ion in a Paul trap [13], the 3-state Potts [14] and Heisenberg [15] anti-ferromagnetic models, scattering [16] and gauge theories [17], molecular adsorption [18], black hole physics [19], as well as quantum anharmonic [20,21] and double well potentials [21]. Moreover, a series expansion approach has been applied to systems with cubic [22] and quartic [23] nonlinearities, which are directly related to the nonlinear version of the Landau-Zener problem considered herein.

The equations to solve are a semiclassical approximation to the photoassociation problem [24], which describes atomic and molecular condensates not as boson fields but as classical fields. We have

$$i\dot{\alpha} = \frac{\Omega}{\sqrt{2}} \alpha^* \beta, \quad i\dot{\beta} = \frac{\Omega}{\sqrt{2}} \alpha^2 + \delta\beta. \quad (5)$$

Physically, α and β are the probability amplitudes that an atom is part of the atomic or the molecular condensate. Equations (5), in fact, preserve the normalization $|\alpha|^2 + |\beta|^2 = 1$. The Rabi frequency for coherent photoassociation, Ω , may be adjusted by varying the intensity of the driving laser field. The detuning δ , again swept linearly as $\delta = \xi t$, measures the mismatch between the laser frequency and the frequency in a transition in which two atoms are converted into a molecule. It is controlled by tuning the photoassociating laser. Transitions to noncondensate modes [25–27] are neglected under the assumption that the condensate coupling satisfies $\Omega \ll \hbar\rho^{2/3}/m$ [4,5,26], where ρ is the initial atom density and m is the atomic mass.

For a fixed detuning the system (5) has dressed states as well [5], though they behave quite differently from the dressed states of the linear system (1). They are of the form $[\alpha(t), \beta(t)] = [e^{-i\mu t}\bar{\alpha}, e^{-2i\mu t}\bar{\beta}]$, where $\hbar\mu$ is conventionally referred to as chemical potential. Depending on the value of the detuning, there may be as many as three essentially different dressed states, along with a multitude of additional states that can be generated using the symmetry transformations of Eqs. (5). A trivial state M exists for all detunings. It has all atoms as molecules, $\bar{\beta} = 1$, and the chemical potential equals $\mu = \frac{1}{2}\delta$. A second state B exists for $\delta/\Omega < \sqrt{2}$ and has $\bar{\alpha} = -1$ at $\delta = -\infty$. For this state $\bar{\beta} \rightarrow 1$ and $\mu \rightarrow \frac{1}{2}\delta$ as $\delta/\Omega \rightarrow 2$, so that at $\delta/\Omega = \sqrt{2}$ the states M and B continuously merge. The third state F exists for $\delta/\Omega > -\sqrt{2}$ and turns into all atoms, $\bar{\alpha} \rightarrow 1$, in the limit $\delta \rightarrow \infty$.

Suppose now that the system starts as all atoms at a large negative detuning, i.e., in state B , and that the detuning is swept slowly through the resonance. One expects adiabatic following, so that the system should stay in state B and eventually turn into state M , into molecules. This actually is the result both in the corresponding quantum case addressed in Refs. [4,5], and from our numerical trials with Eqs. (5). However, the point of resonance where the degree of the transfer of population is mostly determined occurs where the chemical potentials are equal, at $\delta/\Omega = \sqrt{2}$, which is also precisely where the dressed states merge. The situation is mathematically much different from the linear LZ case.

It is therefore gratifying that we have been able to carry out an analysis analogous to the LZ case in our nonlinear system. The (quite involved) technical details will be reported elsewhere, here we only present the key idea. Thus, it is possible to combine from Eqs. (5) a third-order differential equation for the probability for molecules, $P = |\beta|^2$, which reads

$$P''' + \frac{P''}{z} + \left[1 - \frac{1}{4z^2} + \frac{4\lambda}{z} \left(1 - \frac{3}{2}P \right) \right] P' + \frac{\lambda}{z^2} \left(\frac{1}{2} - 2P + \frac{3}{2}P^2 \right) = 0. \quad (6)$$

Here the primes refer to derivatives with respect to the

transformed time variable $z = \xi t^2/2$. The corresponding equation for the linear problem (1) is

$$P''' + \frac{P''}{z} + \left[1 - \frac{1}{4z^2} + \frac{4\lambda}{z}\right] P' + \frac{\lambda}{z^2} [1 - 2P] \equiv \mathcal{D}(\lambda, P) = 0, \quad (7)$$

where $\mathcal{D}$ refers to the rule for forming the left-hand side of the differential equations. The solutions to the latter are, of course, known; $P_{LZ}(z)$ of Eqs. (2) is the one with the appropriate initial conditions at $t = -\infty$.

Now, Eq. (6) may be rearranged to read

$$\mathcal{D}(\lambda, 2P) = 3\lambda \left(\frac{4P'P}{z} - \frac{P^2}{z^2} \right). \quad (8)$$

In the limit with $\lambda \ll 1$ the LZ transition probability is small, and so is presumably the solution to Eq. (6). In Eq. (6) the nonlinear terms, the same ones that make the right-hand side of Eq. (8), are compared with terms that are much larger. The nonlinearity makes a “small” perturbation. We are thus lead to formulate the Ansatz

$$P(z) = \frac{1}{2}P_{LZ}(\lambda, z) + \varepsilon P^{(1)}(z) + \varepsilon^2 P^{(2)}(z) + \dots, \quad (9)$$

with $\varepsilon = P_{LZ}^\infty$ may be thought of as the small parameter in a perturbative expansion.

The calculations make use of what is essentially the Green’s function for the linear differential operator acting on P in Eq. (7). It turns out that a formal expansion of the type (9) may be found, and for a small enough λ it even converges to the desired solution of Eq. (6). The leading terms in the expansion at infinite time are

$$P(\infty) = \frac{1}{2}P_{LZ}^\infty \left(1 + \frac{4\lambda}{\pi} P_{LZ}^\infty \right). \quad (10)$$

Equation (6) may also be rearranged as an equation for the atomic state probability $|\alpha|^2 \equiv R = 1 - P$, and in this form reads

$$\mathcal{D}(-\frac{1}{2}\lambda, R) = 3\lambda \left(-\frac{2R'R}{z} + \frac{R^2 - 1/3}{2z^2} \right). \quad (11)$$

In the limit $\lambda \gg 1$ very few atoms will remain, so one is tempted to attempt an Ansatz of the form

$$R(z) = [1 - P_{LZ}(\frac{1}{2}\lambda, z)] + \varepsilon R^{(1)}(z) + \varepsilon^2 R^{(2)}(z) + \dots. \quad (12)$$

This time the formally small parameter is $\varepsilon = R(\infty) = 1 - P_{LZ}^\infty$. Perhaps surprisingly, this Ansatz works, too. By employing essentially the same mathematics as in the case of the expansion (9), one finds a formal expansion that may be shown to converge for large enough λ . The leading terms give

$$P(\infty) = P_{LZ}^\infty \left(\frac{\lambda}{2} \right) \left[1 - \frac{1}{3\pi\lambda} P_{LZ}^\infty \left(\frac{\lambda}{2} \right) \right]. \quad (13)$$

The most notable qualitative difference from the linear case occurs in the limit of slow sweep of the frequency, $\lambda \gg 1$. The probability for no transition, $|\alpha(\infty)|^2 = 1 - P(\infty)$, behaves in the linear case like $\exp(-4\pi\lambda)$, but in the nonlinear case as $(3\pi\lambda)^{-1}$.

We have solved Eqs. (5) also numerically using several methods. An example is shown in Fig. 1(a), where we plot the atomic probability $R = 1 - P = |\alpha|^2$ as a function of the running detuning $\delta = \xi t$. In this example we initially fix a negative detuning that lies outside the range of the graph, and start the system in the corresponding dressed state B . We then slowly increase the rate of the sweep of the detuning until it reaches $\xi = 0.1\Omega^2$, still far to the left of the detuning axis in Fig. 1(a). Upon arrival to the range plotted in Fig. 1(a), the detuning is swept at this constant rate ξ , and the occupation probability is within 10^{-8} of the occupation probability for the dressed state B . At this point adiabaticity still prevails. Especially around $\delta/\Omega \simeq \sqrt{2}$, though, adiabaticity breaks down. To demonstrate, we show in Fig. 1(b) the difference between the actual occupation probability of the atomic state and the prediction from the dressed state B . Typical of both linear and nonlinear numerical solutions, oscillations arise because the system does not perfectly follow the ground state, and filtering out these oscillations accurately is a major tour de force. Nonetheless, it is easy to see the λ^{-1} dependence of the residual atomic probability on the sweep rate numerically.

While these aspects of the computations are somewhat trivial, computations also provide new insights. The linear LZ problem is completely symmetric in the two states. Whether the system starts in state α or β , the probability for the transition to the other state is the same. This cannot hold in the nonlinear case; if $\alpha = 0$ at some time, then by Eqs. (5) it will remain so forevermore. Also, given that the states B and M merge at $\delta = \sqrt{2}\Omega$, one might expect that there is some preponderance to transitions to a molecular state. This does not seem to be the case. In Fig. 1(c) we resort to the same scheme as in Fig. 1(a), except that we start the system in the superposition with $\alpha = \beta = \frac{1}{\sqrt{2}}$. The probabilities oscillate violently around $\delta = 0$, but eventually settle in the neighborhood of half and half again.

We now briefly consider explicit numbers for ^{23}Na and ^{87}Rb condensates. First is the validity of the two-mode model (5), which again hinges on the condition $\Omega \ll \hbar\rho^{2/3}/m$. The condensate coupling is [5]: $\Omega = [(\lambda^3\rho)(I/I_0)]^{1/2}$, where $2\pi\lambda$ is the wavelength of the photoassociating light and I_0 is the characteristic intensity of a photoassociation transition with binding energy $\sim 1\text{ cm}^{-1}$. Given the ^{23}Na (^{87}Rb) characteristic intensity $I_0 = 0.47\text{ W/cm}^2$ ($I_0 = 0.07\text{ W/cm}^2$) [5], and a typical BEC density $\rho = 5 \times 10^{14}\text{ cm}^{-3}$, the two-mode model is valid for $I \ll 50\text{ mW/cm}^2$ ($I \ll 10\text{ mW/cm}^2$). The linear rate dependence for the transition probability should manifest for slow sweep rates, i.e., $\xi \sim 0.01\Omega^2$. For intensities satisfying the above relation, and typical BEC

densities, this translates into a ^{23}Na (^{87}Rb) detuning-sweep rate $\xi \sim 200 \times 2\pi \text{ Hz}$ ($\xi \sim 50 \times 2\pi \text{ Hz}$).

In sum, we have studied a nonlinear variant of the Landau-Zener problem. Though our specific example was about photoassociation of a Bose-Einstein condensate, the problem is generic in classical and bosonic field theories with a cubic nonlinearity. While the basic adiabaticity argument still works, the structure of the adiabatic or “dressed” states in the nonlinear system is much different from its counterpart in standard linear quantum mechanics. When the levels cross slowly, the probability for the failure of adiabaticity turns out to scale linearly with the rate at which the levels cross, as opposed to the exponential behavior of the linear problem. This prediction could be tested with existing ^{23}Na and ^{87}Rb experimental setups.

-
- [1] L. D. Landau, Phys. Z. Sowjetunion **2**, 46 (1932); C. Zener, Proc. R. Soc. London A **137**, 696 (1932).
 - [2] B. W. Shore, *The Theory of Coherent Atomic Excitation*, (Wiley, New York, 1990), Vol. I; E. E. Nikitin and S. Ya. Umanski, *Theory of Slow Atomic Collisions* (Springer-Verlag, Berlin, 1984); N. V. Vitanov and B. M. Garraway, Phys. Rev. A **53**, 4288 (1996), **54**, 5458(E) (1996); R. Lim and M. V. Berry, J. Phys. A **24**, 3255 (1991).
 - [3] F. Dalfovo, S. Giorgini, L. P. Pitaevskii, and S. Stringari, Rev. Mod. Phys. **70**, 463 (1999).
 - [4] J. Javanainen and M. Mackie, Phys. Rev. A **59**, R3186 (1999).
 - [5] M. Kořtrun, M. Mackie, R. Cote, and J. Javanainen, Phys. Rev. A **62**, 063616 (2000).
 - [6] J. L. Roberts et al., Phys. Rev. Lett. **81**, 5109 (1998); S. Inouye et al., Nature **392**, 151 (1998).
 - [7] P. Tommasini, E. Timmermans, M. Hussein, and A. Kerman, LANL e-print cond-mat/9804015; F. A. van Abeelen and B. J. Verhaar, Phys. Rev. Lett. **83** 1550 (1999); V. A. Yurovsky, A. Ben-Reuven, P. S. Julienne, and C. J. Williams, Phys. Rev. A **60**, R765 (1999).
 - [8] D. F. Walls and C. T. Tindle, J. Phys. A **5**, 534 (1972).
 - [9] For semiclassical theory of a Landau-Zener sweep across a Feshbach resonance without nonlinear implications, see F. H. Mies, E. Tiesinga, and P. S. Julienne, Phys. Rev. A **61**, 022721 (2000).
 - [10] For quantum corrections to the standard Landau-Zener formula applied to photodissociation, see V. A. Yurovsky, A. Ben-Reuven, and P. S. Julienne, LANL e-print, cond-mat/0108372.
 - [11] C. Cohen-Tannoudji, B. Diu, and F. Laloë, *Quantum Mechanics* (Wiley, New York, 1977).
 - [12] M. Abramowitz and I. A. Stegun, *Handbook of Mathematical Functions* (Dover, New York, 1965); A. Erdelyi, W. Magnus, F. Oberhettinger, and F.G. Tricomi, *Higher Transcendental Functions* (McGraw-Hill, New York, 1953).
 - [13] M. Feng and K. L. Wang, Phys. Lett. A **197**, 135 (1995); M. Feng, J. H. Wu, and K. L. Wang, Comm. Theor. Phys. **29** 497 (1998); Z. X. Nie, M. Feng, J. M. Li, L. Shi, X. W. Zhu, and K. L. Gao, *ibid.* **36**, 593 (2001).
 - [14] J. Adler, A. Brandt, W. Janke, S. Shmulyian, J. Phys. A **28**, 5117 (1995).
 - [15] A. W. Sandvik, Phys. Rev. B **56**, 11678 (1997); D. K. Yu, Q. Gu, H. T. Wang, and J. L. Shen, Phys. Rev. B **59**, 111 (1999).
 - [16] T. Gorin, J. Phys. A **32**, 2315 (1999); E. Stefanescu, A. Sandulescu, and W. Scheid, Int. J. Mod. Phys. E **9**, 17 (2000).
 - [17] E. Fahri, V. V. Khoze, K. Rajagopal, and R. Singleton, Phys. Rev. D **50**, 4162 (1994); F. Niedermayer, M. Niedermayer, and P. Weisz, *ibid.* **56**, 2555 (1997); J. M. F. Labastida and E. Perez, J. Math. Phys. **41**, 2658 (2000).
 - [18] D. Poland, J. Chem. Phys. **110**, 6530 (1999); V. Pouthier, J. C. Light, and C. Giradet, *ibid.* **114**, 4955 (2001); C. Fusco, P. Gallo, A. Petri, and M. Rovere, *ibid.* **114**, 7563 (2001).
 - [19] L. M. Burko, Phys. Rev. D **5902**, 4011 (1999); S. Q. Wu and X. Cai, J. Math. Phys. **40**, 4538 (1999).
 - [20] C. M. Bender and L. M. A. Bettencourt, Phys. Rev. Lett. **77**, 4114 (1996); H. Kleinert, S. Thoms, and W. Janke, Phys. Rev. A **55**, 915 (1997); P. B. Kahn and Y. Zarmi, J. Math. Phys. **40**, 4658 (1999).
 - [21] T. Hatsuda, T. Kunihiro, T. Tanaka, Phys. Rev. Lett. **78**, 3229 (1997).
 - [22] S. Chaturvedi and P. D. Drummond, Eur. Phys. J. B **8**, 251 (1999); S. De Palo, C. Castellani, C. Di Castro, and B. K. Chakraverty, Phys. Rev. B **60**, 564 (1999); A. E. Lobo, E. N. Tsoy, C. M. de Sterke, J. Appl. Phys. **90**, 3762 (2001).
 - [23] C. Pare, Phys. Rev. E **54**, 846 (1996); T. A. Laine and A. T. Friberg, *ibid.* **61**, 7098 (2000).
 - [24] We take the semiclassical equations from Ref. [5] and scale the amplitude of the molecular field by $\sqrt{2}$.
 - [25] K. Goral, M. Gadj, and K. Rzazewski, Phys. Rev. Lett. **86**, 1397 (2001); M. Holland, J. Park, and R. Walser, Phys. Rev. Lett. **86**, 1915 (2001).
 - [26] J. Javanainen and M. Mackie, Phys. Rev. Lett. **88**, 090403 (2002).
 - [27] E. A. Donley, N. R. Claussen, S. T. Thompson, and C. E. Wieman, Nature **417**, 529 (2002); S. J. J. M. F. Kokkelmans and M. Holland, cond-mat/0204504; M. Mackie, K.-A. Suominen, and J. Javanainen, cond-mat/0205535.




FIG. 1. (a) Probability of remaining in the atomic condensate $R = 1 - P$ as a function of running detuning $\delta = \xi t$, with $\xi = 0.01 \Omega^2$. The system is started at a large negative time in the dressed eigenstate B , and arrives adiabatically from the left to the range of δ shown in the figure. (b) Difference between the atomic probability R and the adiabatic probability R_B of the dressed state B for the same data as in panel (a). We set $R_B = 0$ for $\delta > \sqrt{2}\Omega$, when the state B has merged with the all-molecules state M . (c) Same as panel (a), except that the system was started in a superposition with $\alpha = \beta = 1/\sqrt{2}$.